




88047187

DEPARTMENT OF THE INTERIOR

POCS REFERENCE PAPER NO. 53-4

Recovery Of Benthic Marine Populations
Along The Pacific Coast Of The United States
Following Natural And Man-made Disturbances
Including Pertinent Life History Information



Prepared by the Bureau of Land Management, Pacific OCS Office
1340 West 6th. Street, Los Angeles, CA 90017

APRIL 1980

TD
195
.P4
P422
1980

BUREAU OF LAND MANAGEMENT
Library
Denver Service Center

BUREAU OF LAND MANAGEMENT
Library
Denver Service Center

BLM LIBRARY
RS 150A BLDG. 50
DENVER FEDERAL CENTER
P.O. BOX 25047
DENVER, CO 80225

IND 88047187

TO
195
P4
P422
1980

RECOVERY OF BENTHIC MARINE POPULATIONS
ALONG THE PACIFIC COAST OF THE UNITED STATES
FOLLOWING NATURAL AND MAN-MADE DISTURBANCES
INCLUDING PERTINENT LIFE HISTORY INFORMATION

By

Lynnette L. Vesco and Robert M. Gillard

POCS Reference Paper No. 53-4
Pacific Outer Continental Shelf Office
Bureau of Land Management
U.S. Department of the Interior

April 1980

BUREAU OF LAND MANAGEMENT.

Library
Denver Service Center

BLM LIBRARY
RS 150A BLDG. 50
DENVER FEDERAL CENTER
P.O. BOX 25047
DENVER, CO 80225

Intertidal and subtidal benthic communities are not static but frequently change in response to natural (e.g., predation and storms) and man-made (e.g., pollution and experimental clearance) disturbances. Studies conducted along the Pacific coast of California, Oregon, and Washington, indicate that the time needed for benthic marine populations to recover from these disturbances ranges from less than 1 month to over 10 years (Table I). Generally, algal populations recover much more rapidly than invertebrate populations. For a community or association to completely recover, all populations, or at least all of the major populations, must become reestablished at pre-disturbance abundances. This typically requires at least 2 years, and often much longer.

It is difficult to determine the precise time required for a population or community to recover because many factors are influential. Therefore, the recovery times given in Table I should be considered approximations. One very important factor that affects recovery times is the severity of the disturbance. For example, if only the erect thallus of the red alga Endocladia muricata is removed, recovery will occur within 6 months (Glynn, 1965) because the alga can regrow from the holdfast; however, if the holdfast is also destroyed, recovery takes over 27 months (Northcraft, 1948) since new spore colonization is required. Furthermore, the more severe the disturbance in terms of the numbers of species and individuals killed or injured, the total area impacted, and the length of the disturbance, the longer it will take a population to recover.

The year or time of year of a disturbance also may influence how long is needed for community recovery. Extensive repopulation by most species will not begin until their setting period which may be up to a year later for those with short reproductive periods. Additionally, reproductive success varies from year to year and in some cases may not be very successful for several years (Paine, 1974). Once established, a species may enhance or inhibit the recolonization of other species. For example, Paine (1977) reported that when an intertidal area in Washington was cleared in the spring while there were many propagules of the green alga Ulva in the water, Ulva encumbered the spatial resource and prevented or delayed invasion of other species. Therefore, this intertidal area took much longer to regain its mature community structure when it was disturbed in the spring (37 months) than when it was cleared in the autumn (20 months).

The time required for a species or an association to recover also varies from one location to the next. Murray and Littler (1979) reported that recovery was generally slower at the northern end of the Southern California Bight than farther south, probably due to differences in water temperature. Therefore, the information in Table I for a particular taxon or assemblage is roughly arranged with the most northern location first followed by sequentially more southerly located areas.

Another factor affecting recovery time is the amount of disturbance to which a population is normally exposed. For example, Murray and Littler

(1978) reported that an area exposed to sewage pollution took only 0.8 months to recover from experimental clearance by scraping and burning, whereas an unpolluted area took more than 30 months.

Finally, recovery time depends on the abundances of species before the disturbance and in nearby undisturbed areas, and the life histories of the species.

All studies of the time required for Pacific coast benthic marine populations to recover from natural and man-made disturbances of which we are aware have been included in Table I. Typically, recovery has been monitored following the experimental clearance of an area by scraping, burning, or exposing new surfaces by chipping or blasting away with dynamite weathered surfaces. Although these studies are certainly valuable, it is important to recognize that scraping does not completely remove crustose forms, burning may leave chemical residues, and exposing new surfaces by chipping or blasting with dynamite alters the natural rock configuration. Table I also includes 1) recovery studies following contamination by oil, 2) recovery studies following natural disturbances such as predation and storms, and 3) studies concerning the colonization of a new substratum when the time needed for a mature community to become established was indicated.

Recovery times are reported either for an individual species or an entire association. Generally, recovery of an association is much longer than for one species because the former is usually based on the species with the slowest recovery rate. Nearly all of the populations

studied primarily occur on hard surfaces. Most also occur in the intertidal zone.

Recovery times have been determined by a variety of methods (see Table I). Perhaps the most feasible quantitative method for determining recovery of the entire community when controls are available is to compare the percent cover of each species to comparable undisturbed data using a Bray-Curtis similarity coefficient or similar analysis (Cimberg, 1975; Murray and Littler, 1978, 1979). Recovery of a population can be quantitatively determined by comparing percent cover data to pre-disturbance data. In comparing cover values by these methods there is no assurance that the typical size or age class distribution has been reestablished since many small individuals could cover the same area. However, this limitation also exists and is probably more pronounced with other methods, such as those based on presence/absence data (i.e., time when settlement was first observed) and density data. Therefore, although Murray and Littler (1979) reported recovery for macroinvertebrate populations based on density as well as cover, only the recovery times based on cover are included in Table I. In extracting recovery times from the tables presented by Murray and Littler, recovery was considered complete when the percent cover was approximately equal to or greater than the pre-disturbance value. Furthermore, only recovery times of the most abundant species from each of their study areas are shown in Table I because recovery of rare species can be easily affected by a few chance settlements and, therefore, could vary greatly each time it is measured.

Since many populations require a long period to recover, complete recovery was not always observed. Occasionally, the recovery times were estimated. If recovery was not observed or predicted, the time was preceded by a "greater than" sign (>) in Table I to indicate a longer period than observed. A population with a >24 month recovery time may actually recover in 25 months or may require over 10 years. Also, a population with a >12 month recovery time will not necessarily recover faster than one with a >24 month recovery time. Recovery of the first population was merely monitored for a shorter period. A "less than or equal to" sign (<=) signifies recovery occurred by the time indicated, but the last monitoring was not for a long period of time (usually 12 months) so it may have occurred much more rapidly.

Several aspects of the life histories of benthic marine invertebrates greatly influence and can be used to predict recovery times. As noted previously, extensive repopulation by most species will not begin until their setting period which may be up to a year later for those with short reproductive periods. After a new generation has settled, it must reach sexual maturity before it can be assumed that a population will persist in an area. To be completely recovered, a population must reestablish its pre-disturbance size or age class structure which typically includes at least a few individuals near their maximum life expectancies.

Tables II and III provide a summary of information on the life histories of Pacific coast benthic marine invertebrates from rocky and soft bottom

habitats, respectively. In obtaining this information the emphasis was placed on rocky intertidal species, but information on other species was included when obtained. As suggested by Giese and Pearse (1974), spawning season was only used to indicate the period when gametes are released, and was distinguished from the brooding, egg-laying and larval release periods. However, earlier authors often did not distinguish these periods, so determining the proper placement of information was sometimes difficult.

Marine invertebrates spawn during different seasons depending on the species. Those that have been thoroughly studied seem to spawn all year or several times per year rather than in distinct annual cycles. However, reproduction in all species is incompletely known and additional information is needed. For example, even though a species may not have a distinct annual cycle through its entire range, populations or individuals in isolated areas could. This assumption is based on the observation that the spawning season of some species seems to depend on geographic location. Populations in Washington sometimes have spawning seasons which do not overlap at all with populations to the south. Occasionally, spawning may merely start later in more northern areas, perhaps in response to an increase in water temperature to some critical value. Similarly, mating seasons also depend on the species, and part of the variation within a species may be attributable to geographic location. Additionally, spawning or mating seasons may vary during succeeding years.

Since reproduction may vary with geographic location, the life history information provided in Tables II and III has been arranged with the most northern area first. Whenever the location was uncertain, it was listed as "Pacific Coast". These references were probably not the original studies. When information provided by one author for a given area encompasses all of the information from others, the data were pooled on one line and all authors referenced. This is particularly prevalent at Monterey Bay where earlier studies were only conducted during the summer months. However, if the findings conflict, the data are provided on separate lines for each author.

Sexual maturity is reached rapidly, usually by 2 years, for most invertebrates. However, some commercially and ecologically valuable taxa such as the spiny lobster Panulirus interruptus and the red abalone Haliotis rufescens do not reach sexual maturity for about 6 years.

The maximum life expectancy of most invertebrates is about 10 years or less. However, several important species live much longer. For example, the sea anemone Anthopleura elegantissima may live 80+ years, chitons live 20-25+ years, the starfish Pisaster ochraceus lives 34 years, the worm Urechis caupo lives about 25 years, and the pismo clam Tivela stultorum lives from 10 to 53 years.

Several methods have been used to determine spawning times, ages at sexual maturity and maximum life expectancies (see Tables II and III). A description of the methods used to determine spawning times is provided by Giese and Pearse (1974). The value of the methods used to

determine ages at sexual maturity and maximum life expectancies depends on the species and geographic location so the methods sections of the individual publications should be consulted. If a method was not given or was not clear, it was assumed that field observations were used.

In general, interpretation of data was avoided. For example, if the larval period and length were known, the setting period could be surmised. However, unless the author indicated the setting period, it was not given in Tables II or III. Similarly, the setting period of many species probably could be approximated from the raw data of recovery studies listed in Table I. However, this was not done since many types of data can be misinterpreted without a thorough understanding of the study.

Data on the life histories of algae have not been tabulated due to time constraints. Many algae produce spores all year: Sargassum muticum (Nicholson, in press), Gigartina agardhii (West, 1972), Macrocystis pyrifera (Oregon State University, 1971), Gigartina spinosa, Iridaea flaccida, Rhodoglossum affine (Northcraft, 1948), while others reproduce seasonally such as Prionitis lanceolata (Northcraft, 1948). Age at sexual maturity is about 1 year for Macrocystis pyrifera (North, 1971). The life span of some algae is quite short: Egregia menziesii 1 to 2 years (Proctor, 1968), Nereocystis luetkeana 1 to 2 years (Carefoot, 1977; Setchell, 1908), Eisenia arborea 2 years (Foster, 1975). However, articulated coralline algae may live over 10 years (Foster, 1975) and clones of Pterocladia capillacea can persist for at least 40 years

(Dixon, 1973).

In summary, repopulation studies indicate that the time needed for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances ranges from less than 1 month to over 10 years. For a community to completely recover, all populations, or at least all of the major populations, must become reestablished at pre-disturbance abundances. This typically requires at least 2 years, and often much longer. Many factors influence the lengths of these recovery times including the severity of the disturbance, the year or time of year of the disturbance, the geographic location, the amount of disturbance to which a population is normally exposed, the abundances of species before the disturbance and in nearby undisturbed areas, and the life histories of the species. Based on life history information, recovery of most species will begin in less than 1 year, but it cannot be assumed they will persist in an area until sexual maturity is reached. This occurs by 2 years for most species. Complete recovery will not occur until the pre-disturbance size or age class structure is reestablished. Often this requires at least a few individuals near their maximum life expectancies and, therefore, may take 10 years or more. Thus, based both on repopulation and life history information, benthic marine communities will need at least 2 years and often 10 years or more to recover from natural and man-made disturbances. Further recovery and life history studies are necessary to evaluate the effects of these disturbances on marine populations, and to more precisely predict recovery times.

ACKNOWLEDGEMENTS

We would like to thank Marty Golden and Steve Murray for providing valuable comments on the manuscript.

We are extremely grateful to Beverly Monroe for typing the numerous pages of tables.

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum.

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
Macroalgae:					
Chlorophyta (green algae)					
<u>Chaetomorpha linum</u>	Dutch Harbor, San Nicolas Isl., CA ¹¹²	I	Rock	Scraped	3 months ^a
<u>Enteromorpha</u> spp.	Dutch Harbor, San Nicolas Isl., CA ¹¹²	I	Rock	Scraped	9 months ^a
<u>Ulva lobata</u>	Government Point, Santa Barbara County, CA ¹¹² Coal Oil Point, Santa Barbara County, CA ¹¹²	I	Rock	Scraped	3 months ^a 3 months ^a
<u>Ulva taeniata</u>	Government Point, Santa Barbara County, CA ¹¹²	I	Rock	Scraped	3 months ^a
Phaeophyta (brown algae)					
<u>Colpomenia sinuosa/peregrina</u>	Fisherman Cove, Santa Catalina Isl., CA ¹¹²	I	Rock	Scraped	3 months ^a
<u>Egregia menziesii</u>	Whites Point, Los Angeles, CA ¹¹² Cave Canyon, Santa Barbara Isl., CA ¹¹² Fisherman Cove, Santa Catalina Isl., CA ¹¹² Wilson Cove, San Clemente Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a 12 months ^a 6 months ^a >24 months ^a
<u>Eisenia arborea</u>	Fisherman Cove, Santa Catalina Isl., CA ¹¹²	I	Rock	Scraped	<24 months ^a
<u>Fucus distichus</u>	Portage Head, WA ³³	I	Rock	Virgin rock surfaces exposed by blasting with dynamite or area soaked in a mixture of white gas and diesel oil and burned	>2-4 years ^a
<u>Halidrys dioica</u>	Willows Anchorage, Santa Cruz Isl., CA ¹¹² Cave Canyon, Santa Barbara Isl., CA ¹¹² Dutch Harbor, San Nicolas Isl., CA ¹¹² Fisherman Cove, Santa Catalina Isl., CA ¹¹² Wilson Cove, San Clemente Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a >24 months ^a >24 months ^a <24 months ^a <24 months ^a

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum (continued).

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
<u>Phaeophyta</u> (continued)					
<u>Hedophyllum sessile</u>	Waadah Isl., WA ¹²⁴	I	Rock	Algal cover scraped off then rock surface sterilized by burning	20 months if cleared in the autumn ^a ; 37 months if cleared in the spring ^a
<u>Hesperophycus harveyanus</u>	Fisherman Cove, Santa Catalina Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a
<u>Laminaria sinclairii</u>	Government Point, Santa Barbara County, CA ¹¹²	I	Rock	Scraped	12 months ^a
<u>Macrocystis pyrifera</u>	Santa Cruz Isl., CA ⁴⁸	S	New concrete construction blocks	--	1 year ^f
<u>Macrocystis pyrifera</u> algal association	Santa Cruz Isl., CA ⁴⁸	S	New concrete construction blocks	--	5-10 years ^f
<u>Pelvetia fastigiata</u>	Government Point, Santa Barbara County, CA ¹¹²	I	Rock	Scraped	>12 months ^a
<u>Pelvetia fastigiata</u> f. <u>gracilis</u>	Cuyler Harbor, San Miguel Isl., CA ¹¹² Willows Anchorage Santa Cruz Isl., CA ¹¹² Fisherman Cove, Santa Catalina Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a >24 months ^a >24 months ^a
<u>Pseudolithoderma nigra</u>	Fisherman Cove, Santa Catalina Isl., CA ¹¹²	I	Rock	Scraped	3 months ^a
<u>Ralfsiaceae</u>	Whites Point, Los Angeles, CA ¹¹² Dutch Harbor, San Nicolas Isl., CA ¹¹²	I	Rock	Scraped	3 months ^a 3 months ^a
<u>Sargassum agardhianum</u>	Wilson Cove, San Clemente Isl., CA ¹¹²	I	Rock	Scraped	>12 months ^a
<u>Rhodophyta</u> (red algae)					
<u>Bosziella orbigniana</u> ssp. <u>dichotoma</u>	Cave Canyon, Santa Barbara Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a
<u>Ceramium eatonlanum/sinicola</u>	Whites Point, Los Angeles, CA ¹¹²	I	Rock	Scraped	3 months ^a
<u>Ceramium</u> spp./ <u>Centroceras clavulatum</u>	Dutch Harbor, San Nicolas Isl., CA ¹¹²	I	Rock	Scraped	3 months ^a

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum (continued).

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
Rhodophyta (continued) <u>Corallina officinalis</u> var. <u>chilensis</u>	Willows Anchorage	1	Rock	Scraped	12 months ^a
	Santa Cruz Isl., CA112				3 months ^a
	Whites Point, CA112				3 months ^a
	Los Angeles, CA112				3 months ^a
	Cave Canyon				3 months ^a
	Santa Barbara Isl., CA112				3 months ^a
	Fisherman Cove, Santa Catalina Isl., CA112				3 months ^a
	Wilson Cove, San Clemente Isl., CA112				3 months ^a
	Ocean Beach, San Diego County, CA112				3 months ^a
	Government Point, Santa Barbara County, CA112	1	Rock	Scraped	>12 months ^a
<u>Corallina vancouveriensis</u>	Coal Oil Point, Santa Barbara County, CA112				3 months ^a
	Santa Barbara County, CA112				>24 months ^a
	Cuyler Harbor, San Miguel Isl., CA112				9 months ^a
	South Point, Santa Rosa Isl., CA112				12 months ^a
	Whites Point, Los Angeles, CA112				12 months ^a
	Cave Canyon, Santa Barbara Isl., CA112				12 months ^a
	Dutch Harbor, San Nicolas Isl., CA112				12 months ^a
	Ocean Beach, San Diego County, CA112				>24 months ^a
	Cuyler Harbor, San Miguel Isl., CA112	1	Rock	Scraped	12 months ^a
	Willows Anchorage, Santa Cruz Isl., CA112				>24 months ^a
Crustose Corallinaceae	Cave Canyon				12 months ^a
	Santa Barbara Isl., CA112				<24 months ^a
	Fisherman Cove, Santa Catalina Isl., CA112				9 months ^a
	Wilson Cove, San Clemente Isl., CA112				12 months ^a
	Ocean Beach, San Diego County, CA112				>24 months ^a
	Whites Point, Los Angeles, CA112	1	Rock	Scraped	>24 months ^a
	Cave Canyon, Santa Barbara Isl., CA112	1	Rock	Scraped	>24 months ^a
	Ocean Beach, San Diego County, CA112				9 months ^a
	Government Point, Santa Barbara County, CA112	1	Rock	Scraped	>12 months ^a
<u>Cryptopleura corallinara</u>					
<u>Cryptopleura crispa</u>					
<u>Cryptopleura violacea</u>					

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum (continued).

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
<u>Rhodophyta</u> (continued)					
<u>Endocladia muricata</u>	Monterey Bay, CA ⁶⁴	I	Rock	Cropped thallus leaving holdfast	>6 months ^e
	Monterey Peninsula, CA ¹¹⁸			Scraped then chipped with sledge hammer until fresh surface uncovered	>27 months ^h
	South Point, Santa Rosa Isl., CA ¹¹²			Scraped	>12 months ^a
	Willows Anchorage, Santa Cruz Isl., CA ¹¹²			Scraped	>24 months ^a
<u>Castroclonium coulteri</u>	Government Point, Santa Barbara County, CA ¹¹²	I	Rock	Scraped	>12 months ^a
<u>Celidium coulteri/pusillum</u>	Willows Anchorage, Santa Cruz Isl., CA ¹¹²	I	Rock	Scraped	6 months ^a
	Fisherman Cove, Santa Catalina Isl., CA ¹¹²				6 months ^a
<u>Celidium purpurascens/robustum</u>	Fisherman Cove, Santa Catalina Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a
<u>Cigartina canaliculata</u>	Government Point, Santa Barbara County, CA ¹¹²	I	Rock	Scraped	>12 months ^a
	Coal Oil Point, Santa Barbara County, CA ¹¹²				12 months ^a
	Guyler Harbor, San Miguel Isl., CA ¹¹²				9 months ^a
	South Point, Santa Rosa Isl., CA ¹¹²				9 months ^a
	Whites Point, Los Angeles, CA ¹¹²				6 months ^a
	Cave Canyon, Santa Barbara Isl., CA ¹¹²				>24 months ^a
	Dutch Harbor, San Nicolas Isl., CA ¹¹²				3 months ^a
	Fisherman Cove, Santa Catalina Isl., CA ¹¹²				6 months ^a
	Wilson Cove, San Clemente Isl., CA ¹¹²				12 months ^a
<u>Cigartina leptorhyncos</u>	Whites Point, Los Angeles, CA ¹¹²	I	Rock	Scraped	3 months ^a
<u>Cigartina papillata</u>	Monterey Peninsula, CA ¹¹⁸	I	Rock	Scraped then chipped with sledge hammer until fresh surface uncovered	>1 ¹ year ^h
<u>Cigartina spinosa</u>	Monterey Peninsula, CA ¹¹⁸	I	Rock	Scraped then chipped with sledge hammer until fresh surface uncovered	>6 months ^h
	Cave Canyon, Santa Barbara Isl., CA ¹¹²			Scraped	>24 months ^a

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum (continued).

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
<u>Rhodophyta</u> (continued)					
<u>Iridaea flaccida</u>	Monterey Peninsula, CA ¹¹⁸	I	Rock	Scraped then chipped with sledge hammer until fresh surface uncovered	>3-4 months ^h
<u>Laurencia sinicola</u>	Willows Anchorage, CA ¹¹² Santa Cruz Isl., CA ¹¹²	I	Rock	Scraped	12 months ^a
<u>Lithothrix aspergillum</u>	Whites Point, CA ¹¹² Los Angeles, CA ¹¹²	I	Rock	Scraped	>24 months ^a
<u>Petrocelis middendorffii</u>	Cuyler Harbor, CA ¹¹² San Miguel Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a
<u>Peyssonneliaceae/Hildenbrandiaceae</u>	South Point, CA ¹¹² Santa Rosa Isl., CA ¹¹²	I	Rock	Scraped	3 months ^a
<u>Plocamium cartilagineum</u>	Government Point, CA ¹¹² Santa Barbara County, CA ¹¹²	I	Rock	Scraped	3 months ^a
<u>Porphyra perforata</u>	Dutch Harbor, CA ¹¹² San Nicolas Isl., CA ¹¹²	I	Rock	Scraped	6 months ^a
<u>Prionitis lanceolata</u>	Cuyler Harbor, CA ¹¹² San Miguel Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a
<u>Pterocladia capillacea</u>	Fisherman Cove, CA ¹¹² Santa Catalina Isl., CA ¹¹² Wilson Cove, CA ¹¹² San Clemente Isl., CA ¹¹²	I	Rock	Scraped	3 months ^a 12 months ^a
<u>Rhodoglossum affine</u>	Monterey Peninsula, CA ¹¹⁸	I	Rock	Scraped then chipped with sledge hammer until fresh surface uncovered	>3-4 months ^h
<u>Rhodomela larix</u>	Government Point, CA ¹¹² Santa Barbara County, CA ¹¹²	I	Rock	Scraped	>12 months ^a
<u>Rhodophycean turf (assemblage of small red algae)</u>	Coal Oil Point, CA ¹¹² Santa Barbara County, CA ¹¹²	I	Rock	Scraped	6 months ^a
<u>Smithora naiadum</u>	Government Point, CA ¹¹² Santa Barbara County, CA ¹¹² Dutch Harbor, CA ¹¹² San Nicolas Isl., CA ¹¹²	I	Rock	Scraped	>12 months ^a >24 months ^a
Plants:					
Spermatophytina (seed plants)					
<u>Phyllospadix scouleri</u>	South Point, CA ¹¹² Santa Rosa Isl., CA ¹¹² Dutch Harbor, CA ¹¹² San Nicolas Isl., CA ¹¹²	I	Rock	Scraped	>12 months ^a <24 months ^a

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum (continued).

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
<u>Spermatophytina</u> (continued)					
<u>Phyllospadix torreyi</u>	Government Point, Santa Barbara County, CA112 Coal Oil Point Santa Barbara County, CA112 Wilson Cove, San Clemente Isl., CA112 Ocean Beach, San Diego County, CA112	I	Rock	Scraped	>12 months ^a 12 months ^a >24 months ^a 12 months ^a
Macroinvertebrates:					
Cnidaria					
Anthozoa (corals, sea anemones)					
<u>Anthopleura elegantissima</u>	Government Point, Santa Barbara County, CA112 Coal Oil Point, Santa Barbara County, CA112 Willows Anchorage, Santa Cruz Isl., CA112 Dutch Harbor, San Nicolas Isl., CA112	I	Rock	Scraped	>12 months ^a >24 months ^a <24 months ^a >24 months ^a
Annelida					
Polychaeta (segmented worms)					
<u>Dodecaceria fewkesi</u>	Cuyler Harbor, San Miguel Isl., CA112	I	Rock	Scraped	>24 months ^a
<u>Phragmatopoma californica</u>	Cuyler Harbor, San Miguel Isl., CA112 South Point, Santa Rosa Isl., CA112 Dutch Harbor, San Nicolas Isl., CA112	I	Rock	Scraped	9 months ^a 9 months ^a 6 months ^a
Arthropoda					
Crustacea (lobsters, crabs, shrimps, barnacles)					
<u>Balanus glandula</u>	Goleta Point, near Santa Barbara, CA144	I	Rock	Contaminated by crude oil due to Platform A blowout	>16 months on exposed surfaces ^h >2 months on sheltered surfaces ^h
	South Point, Santa Rosa Isl., CA112			Scraped	>12 months ^a

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum (continued).

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
Arthropoda - Crustacea (continued)					
<u>Balanus glandula</u> association	Near Trinidad, Humboldt County, CA ²¹	I	Rock	Cleared with hammer, chisel and metal brushes	4.8 years ^b
<u>Balanus improvisus</u>	Fruitvald Avenue Bridge, on Oakland Estuary, CA ⁶⁸ San Francisco Bay, CA ⁶⁸	S	New wooden panels	--	>2.5 months ^d
<u>Chthamalus dalli</u> association	Near Trinidad, Humboldt County, CA ²¹	I	Rock which is periodically buried by sand	Cleared with hammer, chisel and metal brushes	2 months ^j
<u>Chthamalus fissus</u>	Goleta Point, near Santa Barbara, CA ¹⁴⁴	I	Rock	Contaminated by crude oil due to Platform A blowout	>4-6 months ^h
<u>Chthamalus fissus/dalli</u>	Government Point, Santa Barbara County, CA ¹¹² Cuyler Harbor, San Miguel Isl., CA ¹¹² South Point, Santa Rosa Isl., CA ¹¹² Willows Anchorage, Santa Cruz Isl., CA ¹¹² Cave Canyon, Santa Barbara Isl., CA ¹¹² Dutch Harbor, San Nicolas Isl., CA ¹¹² Fisherman Cove, Santa Catalina Isl., CA ¹¹² Wilson Cove, Wilson Cove, San Clemente Isl., CA ¹¹² Ocean Beach, San Diego County, CA ¹¹²	I	Rock	Scraped	>12 months ^a 3 months ^a >12 months ^a >24 months ^a >24 months ^a >24 months ^a >24 months ^a >24 months ^a >24 months ^a
<u>Pachygrapsus crassipes</u>	Duxbury Reef, N. of San Francisco, CA ¹⁶	I	Rock	Contaminated by Bunker C oil from tanker spill	>3 years ^c
<u>Tetracella squamosa rubescens</u>	Fisherman Cove, Santa Catalina Isl., CA ¹¹² Wilson Cove, San Clemente Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a 3 months ^a
Mollusca					
Gastropoda (snails, limpets, abalones)					
<u>Collisella digitatis</u> association	Near Trinidad, Humboldt County, CA ²¹	I	Rock	Cleared with hammer, chisel and metal brushes	<3.25 years ^j
<u>Dendropoma</u> spp. <u>Petaicoconchus montereyensis</u>	Fisherman Cove, Santa Catalina Isl., CA ¹¹²	I	Rock	Scraped	>24 months ^a

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum (continued).

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
Mollusca - Gastropoda (continued) <u>Littorina planaxis</u>	Duxbury Reef, N. of San Francisco, CA ¹⁶	I	Rock	Contaminated by Bunker C oil from tanker spill	>3 years ^c
Mollusca					
Bivalvia (oysters, clams, mussels)					
<u>Chama arcana</u>	Cave Canyon, Santa Barbara Isl., CA ¹²	I	Rock	Scraped	>24 months ^a
<u>Mytilus californianus</u>	Portage Head, WA ³³	I	Rock	Virgin rock surfaces exposed by blasting with dynamite or area soaked in a mixture of white gas and diesel oil and burned	"Many years" ^a
	Oregon (cliff facing NNE) ¹⁴			"Cleared" probably by scraping	>5-6 years ^j
	Oregon (cliff facing SW) ¹⁴			"Cleared" probably by scraping	>8 years ^j
	Duxbury Reef, N. of San Francisco, CA ¹⁸			"Cleaned off" probably by scraping	>10 years ^j
	Monterey Bay, CA ⁸²			Scraped then brushed with a steel brush	>2.5 years ^j
	Cuyler Harbor, San Miguel Isl., CA ¹²			Scraped	>24 months ^a
	South Point, Santa Rosa Isl., CA ¹²			Scraped	>12 months ^a
	Willows Anchorage, Santa Cruz Isl., CA ¹²			Scraped	>24 months ^a
	Whites Point Los Angeles, CA ¹²			Scraped	>24 months ^a
	Corona del Mar, CA ⁹⁸			Predation by the star- fish <u>Pisaster ochraceus</u>	>5-10 years ^j
	Ocean Beach, San Diego County, CA ¹² La Jolla, CA ²⁵			Scraped	>24 months ^a
				Completely cleared by storms	>4 years ^h
<u>Mytilus californianus</u> - <u>Balanus cariosus</u> association	Near Trinidad, Humboldt County, CA ²¹	I	Rock	Cleared with hammer, chisel and metal brushes	9.3 years ^b

Table 1. Time required for benthic marine populations along the Pacific coast of the United States to recover from natural and man-made disturbances or to become established on a new substratum (continued).

Taxon or Association	Geographic Location	Intertidal (I) or Subtidal (S)	Substratum	Type of Disturbance (if any)	Recovery Time
Mollusca - Bivalvia (continued)					
<u>Mytilus edulis</u>	Fruitvald Avenue Bridge on Oakland Estuary, CA ⁶⁸ San Francisco Bay, CA ⁶⁸ Amitos Bay, Long Beach, CA ¹²⁹	S	New wooden panels	--	>5 months ^d
<u>Mytilus edulis</u> - <u>Ulva lobata</u> association	Amitos Bay marina, Long Beach, CA ¹²⁸	S	Boat dock floats New float	Scraped --	2-7 months ^h <1 year ^j
Echinodermata					
Echinodea (sand dollars, sea urchins)					
<u>Strongylocentrotus purpuratus</u>	Willows Anchorage, CA ¹¹² Santa Cruz Isl., CA ¹¹²	I	Rock	Scraped	<24 months ^a
Miscellaneous Associations					
<u>Odonthalia</u> and <u>Iridaea-Redophyllum-Lithothamnion</u> association	Oregon ¹⁴	I	Rock	Cleared probably by scraping	2 years ^j
Combination of 18 species	Duxbury Reef, N. of San Francisco, CA ¹⁷	I	Rock	Contaminated by Bunker C oil from tanker spill	>5 years ^c
"Luxurious fouling growth" (invertebrates)	Monterey Harbor, CA ⁷⁰	I & S	New creosoted wooden piles	--	5-10 years ^j
Sandy beach assemblage	Cat Harbor, Santa Catalina Isl., CA ¹⁴⁵	I	Sandy beach	Contaminated by wet sticky tar	>2 years ^l
<u>Gigartina canaliculata</u> / <u>Corallina officinalis</u> var. chilensis association (unpolluted area)	Wilson Cove, San Clemente Isl., CA ¹¹¹	I	Rock	Scraped, wire brushed, scrubbed, then burned with propane torch	>30 months ^{b,g}
Blue-green algae/ <u>Ulva</u> <u>californica</u> association (polluted area)	Wilson Cove, San Clemente Isl., CA ¹¹¹	I	Rock	Scraped, wire brushed, scrubbed, then burned with propane torch	0.8 months ^{b,g}
"Plant and animal" association	110 miles south of U.S. border in Baja California ¹⁰⁷	I & S	Rock	Contaminated by dark diesel fuel due to grounding of the Tampico-Marú	3-4 years 90% of species restored but some species still not recovered at 12 years ⁱ

LEGEND

Criteria used to determine recovery times:

- a = percent cover data
- b = Bray-Curtis similarity coefficient using percent cover data
- c = density data
- d = size data
- e = dry weight biomass data
- f = growth rate and total length
- g = diversity
- h = time until initial or sizable settlement
- i = distribution or abundance of species
- j = general field observations

Numbers identify the references

Table II. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata.

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Setting Period	Age at Sexual Maturity	Maximum Life Expectancy
Cnidaria Hydrozoa (hydroids, hydrocorals, siphonophores)											
<u>Allopora californica</u>	Central California ¹²¹	S	Males: June-July ^b				Oct-Dec				
<u>Obelia longissima</u>	Pacific Coast ^{99,130}	S						Summer		<1 month ^c	
	Elkhorn Slough, Monterey County, CA ⁹⁷						Aug & Jan				
Cnidaria Anthozoa (corals, sea anemones)											
<u>Anthopleura elegantissima</u>	Pacific Coast ¹³⁰	I									80+ years ^d
	Central California ⁷²										
	San Francisco area, CA ⁴⁷										
	Elkhorn Slough, Monterey County, CA ⁹⁷	I									
Anthopleura xanthogrammica											
Arthropoda Crustacea (lobsters, crabs, shrimps, barnacles)											
<u>Balanus aquila</u>	Monterey Harbor, CA ⁷⁰	I & S							Jan-Mar ⁺		
<u>Balanus cariosus</u>	San Juan Isl., WA ³⁰	I					Spring				
	Near Trinidad, Humboldt County, CA ²¹								Apr ⁺		
<u>Balanus crenatus</u>	Humboldt Bay, CA ⁹⁴	I & S							All year (Peak late summer to early fall)		
	San Francisco Bay Slough, CA ⁸⁰										
	Monterey Bay, CA ¹³⁹										9 months ^f

Table II. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Setting Period	Age at Sexual Maturity	Maximum Life Expectancy
Arthropoda - Crustacea (continued)											
<u>Balanus crenatus</u> (continued)	Monterey Harbor, CA ⁷⁰						Apr ⁺		All year	≥18 months ^c	
<u>Balanus glandula</u>	Vancouver Isl., Canada ³	I		Dec-Jul (Peak) Dec-Jan	Jan-May (Peak Feb) & Aug-Sept Length: 2 months		Mar-May (Peak) Mar-Apr & Aug- Sept	Length: 1 month	Spring		
	San Juan Isl., WA ³⁰						Summer & autumn		May-Oct (Peak May- Aug)	1 year ^f	2 years with predators; >10 years without predators ^f
	Near Trinidad, Humboldt County, CA ²¹								(Peak May & Dec)		
	San Francisco Bay Slough, CA ⁸⁰					(Breeding season discontinued after June)					
	Monterey Bay, CA ⁶⁴								Large part of the year (peak fall and winter)		
	Monterey Bay, CA ^{70,139}									≥1 year ^f	
	Morro Bay, CA ⁸⁸				Dec-May & fall Length: 1 month					6 months ^c	
	Santa Barbara, CA ³⁰										
	Santa Barbara, CA ¹⁴⁴				Jan-June		Jan-Feb & Apr-May		Jan-June		
	La Jolla, CA ³			Oct-Mar (Peak) Oct-Dec	Oct-May (Peak Dec)		(Peak Jan)		Jan-Mar ⁺		
<u>Balanus improvisus</u>	Oakland, CA ⁶⁸	I & S							Mar-Oct (Peak Apr & Sept)		

Table 11. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
Arthropoda - Crustacea (continued)											
<u>Hemigrapsus nudus</u>	Puget Sound, WA ¹³⁰	I			Early summer						
	Monterey, CA ^{12,130}				Oct-May						
<u>Hemigrapsus oregonensis</u>	Monterey Bay, CA ⁸⁴	I			May ⁺						
<u>Ligia occidentalis</u>	Monterey Bay, CA ⁸⁴	I			May-June ⁺						
<u>Pachycheles pubescens</u>	Central Oregon ⁶⁷	I			May-June ⁺						
<u>Pachycheles rudis</u>	Central Oregon ⁶⁷	I			May-June ⁺						
	Elkhorn Slough, Monterey County, CA ⁹⁷				Aug						
	Ocean Beach, San Diego County, CA ⁹⁷				Mar-Aug						
	Ensenada, Baja California ⁹⁷				Dec						
<u>Pachygrapsus crassipes</u>	Pacific Coast ⁸⁵	I			Apr-Sept (Peak Aug)					<1 year ^{C,1}	
	Monterey Bay, CA ^{12,64,84}				Mar-Aug						
<u>Pagurus granosimanus</u>	Monterey Bay, CA ⁸⁴	I			Feb, Apr-May ⁺						
<u>Pagurus hirsutiusculus</u>	Elkhorn Slough, Monterey County, CA ⁹⁷	I			July ⁺			July-Aug			
<u>Pagurus samuelis</u>	Monterey Bay, CA ⁸⁴	I			Feb, Apr-May						
	Monterey Bay, CA ⁹⁷				Aug ⁺			Mar ⁺			
	Monterey Bay, CA ⁷⁵				(Peak spring)				Length: 1 month		3-4 years ^C
	Southern California ⁷⁵				Jan-Oct						
<u>Panulirus interruptus</u>	Pacific Coast ⁹⁹	I & S			<70 days						
	California ^{55,81,154}				Jan-Apr				Length: <2 years	>6 years ^C	>10-11 years ^C
					May-June				Length: 10 weeks		
	California ¹³⁷				(Peak June)			Mar-Aug	Length: 7-8 months		
	California ³⁴				Mar-July				Length: >3-6 months		

Table 11. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
Arthropoda - Crustacea (continued)											
<u>Panulirus interruptus</u> (continued)	San Clemente Isl., CA ¹⁰⁸				May-Aug (Peak June)						>10-11 years ^c
	San Diego, CA ¹³⁵							May-Sept ⁺			
	Southern California and Baja California ⁹²							June-Dec (Peak July-Oct) Length: 7.75 months			
<u>Petrolisthes cinctipes</u>	Pacific Coast ¹³⁰	I			Mar., May- June						
	Central Oregon ⁶⁷				May-June ⁺						
	Elkhorn Slough, Monterey County, CA ⁹⁷				Jan-June						
	Monterey Bay, CA ^{12,84}				All year						
<u>Petrolisthes eriomerus</u>	Central Oregon ⁶⁷	I			May-June ⁺						
<u>Pollicipes polymerus</u>	Monterey Bay, CA ^{86,91}	I			May-Dec Length: 30 days					>6 months ^c	
	Santa Barbara, CA ¹⁴⁴				All year (Peak Feb)						
	Point Dume, Near Santa Monica, CA ⁴									5 years ^f	20 years ^f
<u>Pugettia producta</u>	Elkhorn Slough, Monterey County, CA ⁹⁷	I & S			Feb., Mar, July						
	Monterey Bay, CA ^{12,84}				All year						
<u>Tetraclita squamosa rubescens</u>	Monterey Harbor, CA ⁷⁰	I							July-Dec		
	Morro Bay, CA ⁸⁸				June-Sept Length: 1.5 months					2 years ^c	
Mollusca											
Polyplacophora (chitons)											
<u>Cryptochiton stelleri</u>	Friday Harbor, WA ¹⁰⁰	I & S									25 ⁺ years ^g
	Oregon ¹²⁵		June-July ^c								20 years ^c
	California ⁷⁸		(Breeds Feb-Mar)								
	Monterey Bay, CA ¹⁴⁷		Mar-May ⁺							2 years ^c	

Table II. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
<i>Mollusca</i> - Polyplacophora (continued)											
<i>Katharina tunicata</i>	Central California ^{62,99}	I	Mar-July (peak July) ^{a,c}					July ⁺			
	Monterey Bay, CA ⁸⁴		May ^d							2 years ^c	
	Monterey Bay, CA ⁷⁸										
<i>Mopalia hindsii</i>	Monterey Harbor, CA ⁶²	I & S	Feb-Mar & Nov-Dec ^a								
<i>Mopalia lignosa</i>	Central California ¹⁴⁹	I & S						Length: 5-7 days			
<i>Mopalia muscosa</i>	Puget Sound, WA ⁹⁹	I	July-Aug ^c					Length: 11-12 days			
	Central California ¹⁴⁹										
	Monterey Bay, CA ⁸⁴		Sept ^d								
	Santa Monica Bay, CA ^{10,109}		Winter-spring ^{a,b}								
	Corona del Mar, Orange County, CA ⁹⁹		Nov ^c								
<i>Nuttallina californica</i>	Pacific Coast ⁹⁹	I									20-25 years ^c
<i>Stenoplax heathiana</i>	Pacific Coast ⁹⁹	I						Length: <1 day			
	Monterey, CA ^{76,84}		May-June ^{c,d}								
	Vancouver, Canada ⁸⁷	I & S	Apr ^{bc}								
	Oregon ⁵							Length: 1 week			
<i>Mollusca</i> Gastropoda (snails, limpets, abalone)											
<i>Acanthina punctulata</i>	Monterey Bay, CA ⁸⁴	I				May-June					
<i>Acanthina spirata</i>	Elkhorn Slough, Monterey County, CA ⁹⁷	I				July					
<i>Acmaea mitra</i>	Central California ⁵⁶	I & S	Dec-Mar ^b								
<i>Aplysia californica</i>	Pacific Coast ¹³⁰	I & S				All year					
	California ⁹⁶					Oct-Mar					
	Elkhorn Slough, Monterey County, CA ^{97,99}			July, Aug, Nov ⁺		Nov ⁺				2 years ^c	2 years ^c

Table II. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Setting Period	Age at Sexual Maturity	Maximum Life Expectancy
Mollusca - Gastropoda (continued)											
<u>Collisella asmi</u>	Central California ⁵⁹	I	Mar-Apr & Sept-Oct ^{b,e}								
<u>Collisella digitalis</u>	Coos Bay, OR ⁴⁹	I									6 years ^f
	Rockaway Beach, San Mateo County, CA ⁵⁸		Jan-May ^b								
	Moss Beach, San Mateo County, CA ⁵⁸		Apr, June-July, Dec ^b								
<u>Collisella limatula</u>	Central California ⁵⁸	I	Sept-Oct ^b								
	Monterey Bay, CA ⁸⁴		Sept ^{+d}								
	Palos Verdes, CA ^{133,134}		Apr-Jan (Peak Sept-Oct) ^c								2.75 years ^{f,h}
<u>Collisella pelta</u>	Rockaway Beach, San Mateo County, CA ⁵⁸	I	Fall, winter spring ^b								
<u>Collisella scabra</u>	Bodega Bay, Sonoma County, CA ¹⁴⁶	I	Jan-Mar & fall ^c						All year (Peak July-Oct)		7-11 years ^{f,h}
	Rockaway Beach, San Mateo County, CA ⁵⁹		3 times 1) late winter -early spring 2) early summer 3) late fall ^b								
	Monterey Bay, CA ⁸⁴		Sept ^{+d}								
	Monterey Bay, CA ⁷⁴										
<u>Collisella strigatella</u>	Coos Bay, OR ⁴⁹	I									4 years ^{f,c}
	Santa Barbara, CA ¹⁴⁴		Apr ^{+c}								
<u>Haliotis cracherodii</u>	Monterey Bay, CA ⁸⁴	I & S	Feb-Mar ^{+c}								
	Monterey Bay, CA ^{11,150}		June-Oct ^{a,b}								
	Southern California ⁹⁵		Late spring, summer, early fall ^a								
	Palos Verdes, CA ³²		Spring-early summer ^{+c}								
									(Peak summer & early fall)		

Table 11. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
<i>Mollusca</i> - Gastropoda (continued)											
<i>Haliotis rufescens</i>	California ^{55,154}	I & S						Length: 1-2 weeks			
	Near Mendocino, CA ⁶³		Spring-early summer ^b								
	Central California ³²		Late spring-early fall (Peak late spring-early summer) ^c					Length: 1-2 weeks			
	Monterey, CA ^{9,79,130,132}		Feb-Apr ^c							6 years ^c	
	Monterey Bay, CA ¹¹		All years ^a								
<i>Littorina planaxis</i>	Central California ¹³⁰	I		Most of the year (Peak spring & summer)							
	Monterey Bay, CA ⁸⁴			Mar ⁺		Apr ⁺		May ⁺			
	Monterey Bay, CA ¹¹³			Apr-May							
<i>Littorina scutulata</i>	Monterey, CA ⁶⁴	I							Jan-May ⁺		
<i>Lottia gigantea</i>	Pacific Coast ⁹⁹	I									10-15 years ^f
	Monterey Bay, CA ⁴⁴		Mid winter ^c								
<i>Mitra</i> idae	Southern California ¹⁵	I & S				July-Aug					
	La Jolla & Pt. Loma, San Diego, CA ¹⁹			Feb ⁺		May-July					
<i>Notoacmea fenestrata</i>	Rockaway Beach, San Mateo County, CA ⁵⁷	I	3 times 1) Nov-Dec 2) Jan 3) Mar ^b								
<i>Notoacmea insessa</i>	Moss Beach, San Mateo County, CA ⁵⁷	I	4-5 times per year ^b								
	Monterey Bay, CA ¹²⁷		Summer ^c					Length: 4 days	Summer & spring (Peak June & Aug-Sept)	<1 year ^{c,h}	1 year ^{c,h}
<i>Notoacmea paleacea</i>	Rockaway Beach, San Mateo County, CA ⁵⁷	I	Apparently several times per year ^b								

Table II. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
Mollusca - Gastropoda (continued)											
<u>Notoacmea perona</u>	Oregon ⁹³	I									6+ years ^{g,h}
	Central California ⁵⁷		Jan-Apr (Peak Mar-Apr) ^b								
<u>Notoacmea scutum</u>	Rockaway Beach, San Mateo County, CA ⁵⁷	I	Fall, winter, spring ^b								
<u>Nucella canaliculata</u>	San Juan Isl., WA ¹⁴⁰	I								2-3 years ^c	
	Tomales Bay, Marin County, CA ⁹⁰				Mar-May						
<u>Nucella emarginata</u>	San Juan Isl., WA ¹⁴⁰	I								1.5 years ^c	
	Bodega Bay, CA ⁹⁰					All year (Peak Nov-Feb)					
	Monterey Bay, CA ⁸⁴					Mar, Apr, June					
<u>Petalocochus montereyensis</u>	Monterey, CA ⁷¹	I	(Reproductive all year)								
<u>Serpulorbis squamigerus</u>	Southern California ⁷¹	I	(Females ripe June-Aug ⁺)								
	San Diego, CA ²⁶								Midsummer		
<u>Tegula brunnea</u>	Oregon ⁷	I & S	Aug ⁺ d								
	Monterey, CA ⁸⁴		Mar ⁺ d								
<u>Tegula funebris</u>	Oregon ⁵⁰	I									25 years ^f
	Oregon ⁵¹		Summer ^c						(Peak Jan-Feb)		14-16 years ^g
	Central California ⁵¹										6-8 years ^g
	Monterey Bay, CA ⁸⁴		Apr ⁺ d								
Mollusca											
<u>Bivalvia</u> (oysters, clams, mussels)											
<u>Hinnites giganteus</u>	Pacific Coast ¹³	S									"Several decades" ^c
	Elkhorn Slough, Monterey County, CA ⁹⁷		Apr ⁺ c								

Table II. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
<u>Echinodermata - Echinoidea (continued)</u>											
<u>Strongylocentrotus franciscanus</u> (continued)	Corona del Mar, Orange County, CA ⁹⁹		Dec-Jan ^c								
	Southern California & Baja California ¹²⁶		Spring to early summer ^{+b}								
<u>Strongylocentrotus purpuratus</u>	Oregon ³⁸	I									>10 years ^{f,h}
	Oregon ^{65,66}		Dec-May (Peak Dec-Mar) ^c							2 years ^c	
	Monterey Bay, CA ^{8,60,84,99}		Dec-May ^{a,c}								
	Southern California & Baja California ¹²⁶		Spring-summer ^b								
<u>Echinodermata Asteroidea (starfish)</u>											
<u>Leptasterias hexactis</u>	San Juan Isl., WA ²⁰	I			Nov-Apr Length: 2 months						
	San Juan Isl., WA ^{105,106}				(Peak Mar)					2 years ^c	10 years ^c
	Monterey Bay, CA ^{83,84,99}				Feb-May ⁺						
<u>Leptasterias pusilla</u>	Monterey, CA ¹³⁸	I			Jan-Apr Length: 17-19 days					1-2 years ^c	>2 years ^c
<u>Patiria miniata</u>	Monterey Bay, CA ^{43,99}	I	Most or all year (Peak June-Aug) ^{a,c}								
	Corona del Mar, Orange County, CA ⁹⁹		Jan-July ^c								
<u>Pisaster giganteus</u>	Monterey Bay, CA ^{43,60}	I & S	Jan-March								
<u>Pisaster ochraceus</u>	Pacific Coast ¹³	I									
	San Juan Isl., WA ^{104,105}		June-Aug (Peak June) ^{a,b}								
	San Juan Isl., WA ¹⁰⁶		Mar-June ^{a,d}							5 years ^c	34 years ^c
	Monterey, CA ^{43,60,69,84}										

Table II. Information on the life histories of Pacific coast benthic marine invertebrates associated with rocky substrata (continued).

Taxon	Geographic Location	I or S		Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period		Larval Period	Setting Period	Age at Sexual Maturity	Maximum Life Expectancy
		S	I										
Chordata Ascidacea (sea squirts)													
<u>Styela montereyensis</u>	Del Mar, Near San Diego, CA 131	S									Summer		<30 months ^c

LEGEND

Criteria used to determine spawning seasons, ages at sexual maturity, or maximum life expectancies:

- a = gonad index
- b = gamete ripeness or gonad turgidity
- c = field observations
- d = laboratory observations with or without artificial inducement
- e = thickness of the gonad
- f = growth rates
- g = "annual" growth lines
- h = size class data
- i = animals maintained in aquaria

Other symbols:

- + = the author indicated that this event may also occur during other months of the year
 - I = intertidal
 - S = subtidal
- Numbers identify the references

Table III. Information on the life histories of Pacific coast benthic marine invertebrates occurring in sand or mud habitats.

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
<i>Echiura</i> (unsegmented worms)											
<i>Urechis caupo</i>	Pacific Coast ^{98,99,130}	S	Spring or early summer ^c					Length: <40 days			>25 years ^c
Arthropoda											
Crustacea (lobsters, crabs, shrimps, barnacles)											
<i>Callinassa californiensis</i>	Pacific Coast ^{55,97,130}	I			All year (Peak June-July)						10-16 years ^c
<i>Cancer gracilis</i>	Elkhorn Slough, Monterey County, CA ⁹⁷	I & S		Nov	July-Aug						
<i>Cancer magister</i>	British Columbia, Canada ¹⁰³	S		Apr-Sep	Oct-Aug			Dec-Sept	(Peak Aug)	3-6 years ^c	8-10 years ^c
	Washington ²⁴			Feb-Dec (Peak May-June)	Nov-Feb			Jan-June	June-Aug ⁺	2 years ^c	
	Oregon ¹⁴⁸				Oct-Mar			Apr-July			
	California ^{55,81,119}			Spring	Late fall-early winter			Winter Length: 3-5 months	May ⁺	1.5-2 years ^c	6 years ^c
	Eureka-Crescent City, CA ³⁷				Jan-Feb ⁺						
	San Francisco, CA ³⁷				Jan ⁺					<3 years ^c	7 years ^f
	San Francisco, CA ¹²²										
	San Francisco, CA ⁷⁷										
	Elkhorn Slough, Monterey County, CA ⁹⁷			Mar	All year						

Table III. Information on the life histories of Pacific coast benthic marine invertebrates occurring in sand or mud habitats (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
Arthropoda - Crustacea (continued)											
<u>Emerita analoga</u>	Pacific Coast ^{99,154}	I		Summer or spring	Length: 4-5 months						
	Coos Bay, OR ⁴²				Aug ⁺						
	California ⁴²				All year						
	Monterey, CA ^{12,60}				Apr-Aug						
	Southern California ^{55,130}			Spring-summer				Length: 4 months	(Peak May-July)	1 year ^c	2 years ^c
	Santa Barbara, CA ⁴²								Winter & spring		
	Santa Barbara area, CA ¹⁴⁴				June ⁺			June ⁺			
	Santa Monica, CA ^{41,42}									Males at 2 months, then become females at 14 months ^c	
	La Jolla, CA ^{40,42}					Summer		Length: 4.5 months	Apr-Aug (Peak May-July)		
	La Jolla, CA ³¹					Feb-Sept				Males 2-3 months; females >6-8 months ^c	
<u>Uca crenulata</u>	Southern California ¹¹⁶	I		June-Sept				Apr-Sept			
Mollusca											
Gastropoda (snails, limpets, abalones)											
<u>Olivella biplicata</u>	California ³⁹	I & S (Reproductive all year)						Length: 10-28 days		1 year ^c	>10 years ^c
	San Diego, CA ¹⁴²									2-5 years ^{c,f}	8-12 years ^{c,f}
Mollusca											
Bivalvia (oysters, clams, mussels)											
<u>Crassostrea gigas</u>	California ^{6,36}	I	Spring & fall (Peak June-July) ^{c,d}								
	Elkhorn Slough, Monterey County, CA ⁹⁷	I & S	Summer ^c								
<u>Macoma nasuta</u>											

Table III. Information on the life histories of Pacific coast benthic marine invertebrates occurring in sand or mud habitats (continued).

Taxon	Geographic Location	1 or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Setting Period	Age at Sexual Maturity	Maximum Life Expectancy
Mollusca - Bivalvia (continued)											
<u>Ostrea lurida</u>	Pacific Coast ^{55,130}	1			Length: 10-17 days			Length: 30-40 days		Males 5 months; females 6 months ^c	
	California ⁶				Length: 10-14 days						
	Monterey Bay, CA ⁹⁷							Spring- summer			
	Newport Bay, CA ⁹⁹						July ^{7c}				
	La Jolla, CA ^{25,26,27,28}						Apr-Oct (Peak June-July)		Apr-Nov	4-6 months ^c	
<u>Protothaca staminea</u>	Pacific Coast ^{55,154}	1									7-10 years ^g
	British Columbia, Canada ^{52,53,130}		Jan-Mar ^{b,c}						Feb ⁺	2-3 years ^c	8 years ^c
<u>Saxidomus giganteus</u>	British Columbia, Canada ^{52,54,55}	1 & S	July-Sept ^b					Sept-Nov Length: 4 weeks		3-4 years ^c	17 years ^g
<u>Saxidomus nuttalli</u>	Pacific Coast ⁹⁹	1 & S									10-15 years ^c
	Pacific Coast ^{55,154}		Spring-fall ^c								4-10 years ^c
<u>Siliqua patula</u>	Pacific Coast ⁸¹	1 & S								3-4 years ^c	
	Alaska ^{55,99,130,154}		July-Aug ^c								17 years ^c
	Washington ^{1,55,99,130,154}		May-June ^c								12 years ^c
	California ^{55,154}		Fall ^c					Length: 8 weeks		2 years ^c	6-8 years ^c
<u>Tivela stultorum</u>	Pacific Coast ^{1,45,46,55,99,154}	1 & S	All year (Peak summer & fall) ^c					Length: "several weeks"		1-2 years ^c	26-35 ⁺ years ^c
	Pacific Coast ¹³⁰		Late summer ^c							2-3 years ^c	53 years ^c
	Pacific Coast ⁸¹									3-4 years ^c	
	Central California ²⁹									2 years ^c	10-25 years ^c
	La Jolla, CA ²⁹									1-2 years ^c	10 years ^c
<u>Tresus capax</u>	Humboldt Bay, CA ^{101,102,151}	1 & S	Winter ^{b,c}						Spring		20 years ^f

Table III. Information on the life histories of Pacific coast benthic marine invertebrates occurring in sand or mud habitats (continued).

Taxon	Geographic Location	I or S	Spawning Season	Mating Season	Brooding Period	Egg-Laying Period	Larval Release Period	Larval Period	Settling Period	Age at Sexual Maturity	Maximum Life Expectancy
Mollusca - Bivalvia (continued)											
<u>Tresus nuttalli</u>	Pacific Coast ⁵⁵	I & S	Spring-fall ^c							2 years ^c	
	Pacific Coast ⁸¹									3-4 years ^c	
	Pacific Coast ¹⁵⁴		Winter-early spring ^c								
	Elkhorn Slough, Monterey County, CA ^{22,99}		All year (Peak Feb-Apr) ^c					Length: 21-30 days	All year (Peak Apr-May)	>2 years ^c	>17 years ^c
<u>Zirfaea pilsbryi</u>	Pacific Coast ^{99,130}	I & S	July ^{+c}								7-8 years ^c
Echinodermata											
Echinodea (sand dollars, sea urchins)											
<u>Dendraster excentricus</u>	Monterey Bay, CA ⁹⁷	I & S	July-Aug ^{+c}								
	San Diego area, CA ¹¹⁵		Early spring- midsummer (Peak spring) ^a								

LEGEND

Criteria used to determine spawning seasons, ages at sexual maturity, or maximum life expectancies:

- a = gonad index
- b = Gamete ripeness or gonad turgidity
- c = field observations
- d = laboratory observations with or without artificial inducement
- e = thickness of the gonad
- f = growth rates
- g = "annual" growth lines
- h = size class data
- i = animals maintained in aquaria

Other symbols:

+ = the author indicated that this event may also occur during other months of the year

I = intertidal

S = subtidal

Numbers identify the references

LITERATURE CITED

1. Amos, M. H. 1966. Commercial clams of the North American Pacific coast. Circular 237, U.S. Dept. of the Interior, Fish and Wildlife Service, Bureau of Commercial Fisheries.
2. Anderson, W. R. and R. F. Ford. 1976. Early development, growth and survival of the yellow crab Cancer anthonyi Rathbun (Decapoda, Brachyura) in the laboratory. Aquaculture 7:267-279.
3. Barnes, H. and M. Barnes. 1956. The general biology of Balanus glandula Darwin. Pac. Sci. 10:415-422.
4. Barnes, H. and E. S. Reese. 1960. The behaviour of the stalked intertidal barnacle Pollicipes polymerus J. B. Sowerby, with special reference to its ecology and distribution. J. Anim. Ecol. 29:169-185.
5. Barnes, J. R. and J. J. Gonor. 1973. The larval settling of the lined chiton Tonicella lineata. Mar. Biol. 20:259-264.
6. Barrett, E. M. 1963. The California oyster industry. Calif. Dept. Fish and Game Fish Bull. 123. 103 pp.
7. Belcik, F. P. 1965. Note on a range extension and observations of spawning in Tegula, a gastropod. Veliger 7:233-234.
8. Bennet, J. and A. C. Giese. 1955. The annual reproductive and nutritional cycles in two western sea urchins. Biol. Bull. 109:226-237.
9. Bonnot, P. 1930. Abalones in California. Calif. Fish and Game 16:15-23.
10. Boolootian, R. A. 1964. On growth, feeding and reproduction in the chiton Mopalia muscosa of Santa Monica Bay. Helgo. Wiss. Meer. 11:186-199.
11. Boolootian, R. A., A. Farmanfarmian and A. C. Giese. 1962. On the reproductive cycle and breeding habits of two western Haliotis. Biol. Bull. 122:183-193.
12. Boolootian, R. A., A. C. Giese, A. Farmanfarmian, and J. S. Tucker. 1959. Reproductive cycles of five west coast crabs. Physiolog. Zool. 32:213-220.
13. Carefoot, T. 1977. Pacific seashores. Univ. Wash. Press: Seattle. 208 pp.

14. Castenholz, R. 1967. Stability and stresses in intertidal populations. pp. 15-18. In T. A. Olson and F. J. Burgess (eds.). Pollution and marine ecology. Interscience publishers: New York. 364 pp.
15. Cate, J. M. 1968. Mating behavior in Mitra idae (Melvill, 1893). Veliger 10:247-252.
16. Chan, G. 1975. A study of the effects of the San Francisco oil spill on marine life. Part II: Recruitment. pp. 457-461. In Proc. 1975 conference on prevention and control of oil pollution. March 25-27. San Francisco, California. Sponsored by A.P.I., E.P.A. and U.S.C.G.
17. Chan, G. 1977. The five-year recruitment of marine life after the 1971 San Francisco oil spill. pp. 543-545. In Proc. 1977 oil spill conference. March 8-10, 1977. New Orleans, Louisiana. Sponsored by A.P.I., E.P.A., and U.S.C.G.
18. Chan, G. L. and A. Molina. 1969. The conservation of marine animals on Duxbury Reef. College of Marin, Kentfield, CA 94904. 56 pp.
19. Chess, J. R. and R. J. Rosenthal. 1971. On the reproductive biology of Mitra idae (Gastropoda: Mitridae). Veliger 14:172-176.
20. Chia, F.-S. 1966. Brooding behavior of a six-rayed starfish, Leptasterias hexactis. Biol. Bull. 130:304-315.
21. Cimberg, R. L. 1975. Zonation, species diversity and redevelopment in the rocky intertidal near Trinidad, Northern California. M.S. Thesis, Humboldt State Univ. 118 pp.
22. Clark, P. J. Nybakken and L. Laurent. 1975. Aspects of the life history of Tresus nuttallii in Elkhorn Slough. Calif. Fish and Game 61:215-227.
23. Clarke, W. D. 1967. Part II: Discussion. pp. 43-50. In T. A. Olson and F. J. Burgess (eds.). Pollution and marine ecology. Interscience Publishers: New York. 364 pp.
24. Cleaver, F. C. 1949. Preliminary results of the coastal crab (Cancer magister) investigation. Wash. State. Dept. Fish., Biol. Report 49-A:47-82.
25. Coe, W. R. 1930. The life cycle of the California oyster (O. lurida). Anat. Rec. 47:359.

26. Coe, W. R. 1932a. Season of attachment and rate of growth of sedentary marine organisms. Bull. Scripps Inst. Oceanogr. Tech. Ser. 3:37-86.
27. Coe, W. R. 1932b. Development of the gonads and the sequence of the sexual phases in the California oyster (Ostrea lurida). Bull. Scripps Inst. Ocean. Tech. Ser. 3:119-144.
28. Coe, W. R. and W. E. Allen. 1937. Growth of sedentary marine organisms on experimental blocks and plates for nine successive years at the pier of the Scripps Institution of Oceanography. Bull. Scripps Inst. Oceanogr. Tech. Ser. 4:101-136.
29. Coe, W. R. and J. E. Fitch. 1950. Population studies, local growth rates and reproduction of the pismo clam. Jour. Mar. Res. 9:188-210.
30. Connell, J. H. 1970. A predatory-prey system in the marine intertidal region. 1. Balanus glandula and several predatory species of Thais. Ecol. Monogr. 40:49-78.
31. Cox, G. W. and G. H. Dudley. 1968. Seasonal pattern of reproduction of the sand crab Emerita analoga in southern California. Ecology 49:746-751.
32. Cox, K. W. 1962. California abalones, family Haliotidae. Calif. Dept. Fish and Game Fish Bull. 118. 133 pp.
33. Dayton, P. K. 1971. Competition, disturbance, and community organization: the provision and subsequent utilization of space in a rocky intertidal community. Ecol. Monogr. 41:351-389.
34. Dees, L. T. 1963. Spiny lobsters. Fishery leaflet 523, U.S. Dept. of the Interior, Fish and Wildlife Service, Bureau of Commercial Fisheries.
35. Dixon, P. S. 1973. Biology of the Rhodophyta. Oliver & Boyd: Edinburgh. 285 pp.
36. Ebert, E. E., A. W. Haseltine and R. O. Kelly. 1974. Seawater system design and operations of the marine culture laboratory, Granite Canyon. Calif. Fish and Game 60:4-14.
37. Ebert, E. E., R. O. Kelly and A. W. Haseltine. 1975. Observations on the larval hatching success of the Dungeness crab, Cancer magister, from the San Francisco and Eureka-Crescent City region. Calif. Dept. Fish and Game, Marine Resources Technical Rept. 29. 14 pp.

38. Ebert, T. A. 1967. Negative growth and longevity in the purple sea urchin Strongylocentrotus purpuratus (Stimpson). Science 157:557-558.
39. Edwards, D. C. 1968. Reproduction in Olivella biplicata. Veliger 10:297-304.
40. Efford, I. E. 1965. Aggregation in the sand crab, Emerita analoga (Stimpson). J. Anim. Ecol. 34:63-75.
41. Efford, I. E. 1967. Neoteny in sand crabs of the genus Emerita (Anomura, Hippidae). Crustaceana 13:81-93.
42. Efford, I. E. 1970. Recruitment to sedentary marine populations as exemplified by the sand crab (Emerita analoga) (Decapoda, Hippidae). Crustaceana 18:293-308.
43. Farmanfarmaian, A., A. C. Giese, R. A. Boolootian, and J. Bennett. 1958. Annual reproductive cycles in four species of west coast starfishes. J. Exp. Zool. 138:355-367.
44. Fisher, W. K. 1904. Anatomy of Lottia gigantea. Zool. Jahrb. Abt. Anat. 20:1-66.
45. Fitch, J. E. 1953. Common marine bivalves of California. Calif. Dept. Fish and Game Fish Bull. 90. 102 pp.
46. Fitch, J. E. 1961. The pismo clam. Calif. Dept. Fish and Game, Marine Resources Leaflet No. 1. 22 pp.
47. Ford, C. E. 1964. Reproduction in the aggregating sea anemone, Anthopleura elegantissima. Pac. Sci. 18:138-145.
48. Foster, M. 1975. Algal succession in a Macrocystis pyrifera forest. Mar. Bio. 32:313-329.
49. Frank, P. W. 1965a. Growth of three species of Acmaea. Veliger 7:201-202.
50. Frank, P. W. 1965b. Shell growth in a natural population of the turban snail, Tegula funebris. Growth 29:395-403.
51. Frank, P. W. 1975. Latitudinal variation in the life history features of the black turban snail Tegula funebris (Prosobranchia: Trochidae). Mar. Biol. 31:181-192.
52. Fraser, C. M. 1929. The spawning and free-swimming larval periods of Saxidomus and Paphia. Trans. Roy. Soc. Can. Ser. 3, Vol. 23, Sect. V:195-198.

53. Fraser, C. M. and G. M. Smith. 1928a. Notes on the ecology of the little-neck clam, Paphia staminea Conrad. Trans. Roy. Soc. Can. Ser. 3, Vol. 22, Sect. V:249-269.
54. Fraser, C. M. and G. M. Smith. 1928b. Notes on the ecology of the butter clam, Saxidomus giganteus. Trans. Roy. Soc. Can. Ser. 3, Vol. 22, Sect. V:271-286.
55. Frey, H. W. 1971. California's living marine resources and their utilization. Calif. Dept. Fish and Game. 148 pp.
56. Fritchman, H. K. II. 1961a. A study of the reproductive cycle in California Acmaeidae (Gastropoda). Part I. Veliger 3:57-63.
57. Fritchman, H. K. II. 1961b. A study of the reproductive cycle in California Acmaeidae (Gastropoda). Part II. Veliger 3:95-100.
58. Fritchman, H. K. II. 1961c. A study of the reproductive cycle in California Acmaeidae (Gastropoda). Part III. Veliger 4:41-47.
59. Fritchman, H. K. II. 1962. A study of the reproductive cycle in California Acmaeidae (Gastropoda). Part IV. Veliger 4:134-40.
60. Giese, A. C. 1959. Reproductive cycles of some west coast invertebrates. pp. 625-638. In R. B. Withrow (ed.). Photoperiodism and related phenomena in plants and animals. Am. Ass. Adv. Sci. Washington Publ. No. 55.
61. Giese, A. C. and J. S. Pearse. 1974. Reproduction of marine invertebrates. Vol. I. Acoelomate and pseudocoelomate metazoans. Academic Press: New York. 546 pp.
62. Giese, A. C., J. S. Tucker and R. A. Boolootian. 1959. Annual reproductive cycles of the chitons Katharina tunicata and Mopalia hindsii. Biol. Bull. 117:81-88.
63. Giorgi, A. E. and J. D. DeMartini. 1977. A study of the reproductive biology of the red abalone, Haliotis rufescens Swaison, near Mendocino, California. Calif. Fish and Game 63:80-94.
64. Glynn, P. W. 1965. Community composition, structure, and interrelationships in the marine intertidal Endocladia muricata - Balanus glandula association in Monterey Bay, California. Beaufortia (Zool. Mus., Amsterdam) 12:1-198.
65. Gonor, J. J. 1972. Gonad growth in the sea urchin, Strongylocentrotus purpuratus (Stimpson) (Echinodermata: Echinodea) and the assumptions of gonad index methods. J. Exp. Mar. Biol. Ecol. 10:89-103.

66. Gonor, J. J. 1973. Reproductive cycles in Oregon populations of the echinoid, Strongylocentrotus purpuratus (Stimpson). I. Annual gonad growth and ovarian gametogenic cycles. J. Exp. Mar. Biol. Ecol. 12:45-64.
67. Gonor, S. L. and J. J. Gonor. 1973. Descriptions of the larvae of four North Pacific Porcellanidae (Crustacea: Anomura). Fishery Bulletin 71:189-224.
68. Graham, H. W. and Helen Gay. 1945. Season of attachment and growth of sedentary marine organisms at Oakland, California. Ecology 26:375-386.
69. Greenfield, L. J. 1959. Biochemical and environmental factors involved in the reproductive cycle of the sea star Pisaster ochraceus (Brandt). Ph.D. Thesis, Stanford Univ. 143 pp.
70. Haderlie, E. C. 1974. Growth rates, depth preference and ecological succession of some sessile marine invertebrates in Monterey Harbor. Veliger 17(Suppl.):1-35.
71. Hadfield, M. G. 1966. The reproductive biology of the California vermetid gastropods Serpulorbis squamigerus (Carpenter, 1857) and Petalochuchus montereyensis Dall, 1919. Ph.D. Thesis, Stanford Univ. 174 pp.
72. Hand, C. 1955. The sea anemones of central California. Part II. The endomyarian and mesomyarian anemones. Wasmann J. Biol. 13:37-99.
73. Harger, J. R. E. 1970. Comparisons among growth characteristics of two species of sea mussel, Mytilus edulis and M. californianus. Veliger 13:44-56.
74. Haven, S. B. 1971. Niche differences in the intertidal limpets Acmaea scabra and Acmaea digitalis (Gastropoda) in central California. Veliger 13:231-248.
75. Hays, R. P. 1939. Notes on the life history of the hermit crab Pagurus samuelis. M.A. Thesis, Stanford Univ. 58 pp.
76. Heath, H. 1899. The development of Ischnochiton. Zool. Jahrb. Abt. Anat. 12:567-656.
77. Heath, H. 1902. The breeding habits of Cancer magister. Amer. Nat. 36:501-502.
78. Heath, H. 1905. Breeding habits of chitons of the California coast. Zool. Anz. 29:390-393.

79. Heath, H. 1925. The abalone question. Calif. Fish and Game 11:138-139.
80. Herz, L. E. 1933. The morphology of the later stages of Balanus crenatus Bruguiere. Biol. Bull. 64:432-442.
81. Hester, F. J. 1974. The living marine resources of central and southern California and Baja California. Prepared for Western Oil and Gas Association. 161 pp.
82. Hewatt, W. G. 1935. Ecological succession in the Mytilus californianus habitat, as observed in Monterey Bay, California. Ecology 16:244-251.
83. Hewatt, W. G. 1937. Ecological studies on selected marine intertidal communities of Monterey Bay, California. Amer. Midland Nat. 18:161-206.
84. Hewatt, W. G. 1938. Notes on the breeding seasons of the rocky beach fauna of Monterey Bay, California. Proc. Calif. Acad. Sci. (4)23(19):283-288.
85. Hiatt, R. M. 1948. The biology of the lined shore crab, Pachygrapsus crassipes Randall. Pac. Sci. 2:135-213.
86. Hilgard, G. H. 1960. A study of reproduction in the intertidal barnacle Mitella polymerus, in Monterey Bay, California. Biol. Bull. 119:169-188.
87. Himmelman, J. H. 1975. Phytoplankton as a stimulus for spawning in three marine invertebrates. J. Exp. Mar. Biol. Ecol. 20:199-214.
88. Hines, A. H. 1978. Reproduction in three species of intertidal barnacles from central California USA. Biol. Bull. 154:262-281.
89. Hines, A. H. 1979. Effects of a thermal discharge on reproductive cycles in Mytilus edulis and Mytilus californianus (Mollusca, Bivalvia). Fishery Bulletin 77:498-503.
90. Houston, R. S. 1971. Reproductive biology of Thais emarginata (Deshayes, 1839) and Thais canaliculata (Duclos, 1832). Veliger 13:348-357.
91. Howard, G. K. 1959. Aspects of nutrition and reproduction in the barnacle Mitella polymerus in Monterey Bay, California. M.A. Thesis, Stanford Univ. 105 pp.

92. Johnson, M. W. 1960. Production and distribution of larvae of the spiny lobster Panulirus interruptus (Randall) with records on P. gracilis Streets. Bull. Scripps Inst. Oceanog. 7:413-462.
93. Kenny, R. 1969. Growth characteristics of Acmaea persona Eschscholtz. Veliger 11:336-339.
94. Lambert, T. R. 1973. Seasons of attachment, settlement on various substrates and abundance of colonizing organisms on an artificial reef in Humboldt Bay, California. M.S. Thesis, Humboldt State University. 118 pp.
95. Leighton, D. and R. Boolootian. 1963. Diet and growth in the black abalone, Haliotis cracherodii. Ecology 44:227-238.
96. MacGinitie, G. E. 1934. The egg-laying activities of the sea hare Tethys californicus (Cooper). Biol. Bull. 67:300-303.
97. MacGinitie, G. E. 1935. Ecological aspects of a California marine estuary. Amer. Midl. Nat. 16:629-765.
98. MacGinitie, G. E. 1938. Notes on the natural history of some marine animals. Amer. Midl. Nat. 19:207-219.
99. MacGinitie, G. E. and N. MacGinitie. 1968a. Natural history of marine animals. Second edition. McGraw-Hill: New York. 523 pp.
100. MacGinitie, G. E. and N. MacGinitie. 1968b. Notes on Cryptochiton stelleri (Middendorff, 1846). Veliger 11:59-61.
101. Machell, J. R. 1968. The reproductive cycle of the clam Tresus capax (Gould, 1850), family Mactridae in south Humboldt Bay, California. M.S. Thesis, Humboldt State Univ. (Abstract).
102. Machell, J. R. and J. D. DeMartini. 1971. An annual reproductive cycle of the gaper clam, Tresus capax (Gould), in south Humboldt Bay, California. Calif. Fish and Game 57:274-282.
103. MacKay, D. C. G. 1942. The Pacific edible crab, Cancer magister. Fish. Res. Bd. Can., Bull. 62:1-32.
104. Mauzey, K. 1966. Feeding, behavior and reproductive cycles in Pisaster ochraceus. Biol. Bull. 131:127-144.
105. Menge, B. A. 1974. Effect of wave action and competition on brooding and reproductive effort in the sea star, Leptasterias hexactis. Ecology 55:84-93.

106. Menge, B. A. 1975. Brood or broadcast? The adaptive significance of different reproductive strategies in the two intertidal sea stars Leptasterias hexactis and Pisaster ochraceus. Mar. Biol. 31:87-100.
107. Mitchell, C. T., E. K. Anderson, L. G. Jones and W. J. North. 1970. What oil does to ecology. J. Wat. Pollut. Control Fed. 42:812-818.
108. Mitchell, C. T., C. H. Turner and A. R. Strachan. 1969. Observations on the biology and behavior of the California spiny lobster, Panulirus interruptus (Randall). Calif. Fish and Game 55:121-131.
109. Monroe, H. C. and R. A. Boolootian. 1965. Reproductive biology in the chiton Mopalia muscosa. Bull. So. Calif. Acad. Sci. 64:223-228.
110. Moore, D. R. and D. J. Reish. 1969. Studies on the Mytilus edulis community in Alamitos Bay, California. IV. Seasonal variation in gametes from different regions in the Bay. Veliger 11:250-255.
111. Murray, S. N. and M. M. Littler. 1978. Patterns of algal succession in a perturbed marine intertidal community. J. Phycol. 14:506-512.
112. Murray, S. N. and M. M. Littler. 1979. Experimental studies of the recovery of populations of rocky intertidal macro-organisms following mechanical disturbance. Science Applications, Inc., Tech. Rept. II-2.0 to the BLM. Contract AA550-CT7-44 (Year-III SCOCs Program), La Jolla.
113. Neale, J. R. 1965. Rheotactic responses in the marine mollusk Littorina planaxis Philippi. Veliger 8:7-10.
114. Nicholson, N. in press. The biology of Sargassum muticum (Yendo) Fensholt at Santa Catalina Island, California. Int. Seaweed Symp. 8.
115. Niesen, T. M. 1977. Reproductive cycles in two populations of the Pacific sand dollar Dendraster excentricus. Mar. Biol. 42:365-374.
116. Nishimoto, R. T. 1974. Ecology of Uca crenulata (Lockington) in upper Newport Bay, California. M.A. Thesis, Calif. State Univ., Fullerton. 135 pp.
117. North, W. J. (ed.). 1971. The biology of giant kelp beds (Macrocystis) in California. Beihefte Nova Hedwigia 32:1-600.

118. Northcraft, R. D. 1948. Marine algal colonization on the Monterey Peninsula, California. Amer. J. Bot. 35:396-404.
119. Orcutt, H. G., A. N. Tasto, P. W. Wild, C. W. Hauger, and E. E. Ebert. 1976. Dungeness crab research program report for the year 1976. Dept. Fish and Game, Marine Resources Admin. Report. No. 76-15. 42 pp.
120. Oregon State University. 1971. Oceanography of the nearshore coastal waters of the Pacific Northwest relating to possible pollution. Prep. for U.S. EPA, Water Quality Office Grant No. 16080 EDK, OSU, Corvallis, OR. Vol I, 615 pp.; Vol II, 744 pp.
121. Ostarello, G. L. 1973. Natural history of the hydrocoral Allopora californica Verrill (1866). Biol. Bull. 145:548-564.
122. Pacific Fishery Management Council. 1979. Draft fishery management plan for the Dungeness crab fishery off Washington, Oregon and California. 93 pp.
123. Paine, R. J. 1974. Intertidal community structure: experimental studies on the relationship between a dominant competitor and its principal predator. Oecologia 15:93-120.
124. Paine, R. J. 1977. Controlled manipulations in the marine intertidal zone and their contributions to ecological theory. pp. 245-270. In Changing scenes in natural sciences 1776-1976. Academy of Natural Sciences. Special Publication 12.
125. Palmer, J. B. and P. W. Frank. 1974. Estimates of growth of Cryptochiton stelleri (Middendorff, 1846) in Oregon. Veliger 16:301-304.
126. Pearse, J. S. 1971. Further observations on the reproduction of sea urchins. pp. 108-115. In W. J. North (ed.). Kelp habitat improvement project annual report 1970-1971.
127. Proctor, S. J. 1968. Studies on the stenotopic marine limpet Acmaea insessa (Mollusca: Gastropoda: Prosobranchia) and its algal host Egregia menziesii (Phaeophyta). Ph.D. Thesis, Stanford Univ. 144 pp.
128. Reish, D. J. 1962. A study of succession in recently constructed marine harbors in Southern California. pp. 570-573. In Proc. first national coastal and shallow water research conference. Baltimore, Los Angeles and Tallahassee.
129. Reish, D. J. 1964. Studies on the Mytilus edulis community in Alamitos Bay, California: I. Development and destruction of the community. Veliger 6:124-131.

130. Ricketts, E. F. and J. Calvin. 1968. Between Pacific tides. 4th ed. Revised by J. W. Hedgpeth. Stanford Univ. Press: Stanford. 614 pp.
131. Rosenthal, R., W. D. Clarke and P. K. Dayton. 1974. Ecology and natural history of a stand of giant kelp, Macrocystis pyrifera, off Del Mar, California. Fishery Bulletin 72:670-684.
132. Scofield, N. B. 1930. Conservation laws provide ample protection for abalones. Calif. Fish and Game 16:13-15.
133. Seapy, R. R. 1966. Reproduction and growth in the file limpet, Acmaea limatula Carpenter, 1864. Veliger 8:300-310.
134. Segal, E. 1956. Microgeographic variation as thermal acclimation in an intertidal mollusc. Biol. Bull. 111:129-152.
135. Serfling, S. A. and R. F. Ford. 1975. Ecological studies of the puerulus larval stage of the California spiny lobster Panulirus interruptus. Fishery Bulletin 73:360-377.
136. Setchell, W. A. 1908. Nereocystis and Pelagophycus. Bot. Gaz. 45:125-134.
137. Seymour, G. 1971. Lobster. Dept. Fish and Game, Wildlife Leaflet 81521-800. 2 pp.
138. Smith, R. H. 1971. Reproductive biology of a brooding sea-star, Leptasterias pusilla (Fisher), in the Monterey Bay Region. Ph.D. Thesis, Stanford Univ. 214 pp.
139. Smith, S. V. and E. C. Haderlie. 1969. Growth and longevity of some calcareous fouling organisms, Monterey Bay, California. Pac. Sci. 23:447-451.
140. Spight, T. M. 1975. On a snail's chance of becoming a year old. Oikos 26:9-14.
141. Stohler, R. 1930. Beitrag zur Kenntnis des Geschlechtszyklus von Mytilus californianus Conrad. Zool. Anz. 90:263-268.
142. Stohler, R. 1969. Growth study in Olivella biplicata (Sowerby, 1825). Veliger 11:259-267.
143. Straughan, D. 1971a. The influence of oil and detergents on recolonization in the upper intertidal zone. pp. 437-440. In Proc. joint conf. on prevention and control of oil spills. Sponsored by A.P.I., E.P.A. and U.S.C.G.

144. Straughan, D. 1971b. Breeding and larval settlement of certain intertidal invertebrates in the Santa Barbara Channel following pollution by oil. pp. 223-244. In D. Straughan (ed.). Biological and oceanographical survey of the Santa Barbara Channel oil spill 1969-1970. V. I. Biology and bacteriology. Allan Hancock Foundation, Univ. Southern California. 426 pp.
145. Straughan, D. 1979. Distribution of tar and relationship to changes in intertidal organisms on sandy beaches in Southern California. pp. 591-601. In Proc. 1979 oil spill conference. March 19-22, 1979. Los Angeles, California. Sponsored by A.P.I., E.P.A., and U.S.C.G.
146. Sutherland, J. 1970. Dynamics of high and low populations of the limpet Acmaea scabra (Gould). Ecol. Monogr. 40:169-188.
147. Tucker, J. S. and A. C. Giese. 1962. The reproductive cycles of Cryptochiton stelleri (Middendorff). J. Exper. Zool. 150:33-43.
148. Waldron, K. D. 1958. The fishery and biology of the Dungeness crab (Cancer magister Dana) in Oregon waters. Fish. Comm. Oregon, Contr. No. 24. 43 pp.
149. Watanabe, J. and L. R. Cox. 1975. Spawning behavior and larval development in Mopalia lignosa and Mopalia muscosa (Mollusca: Polyplacophora) in central California. Veliger 18(Supl.):18-27.
150. Webber, H. H. and A. C. Giese. 1969. Reproductive cycles and gametogenesis in the black abalone Haliotis cracherodii (Gastropoda: Prosobranchiata). Mar. Bio. 4:152-159.
151. Wendell, F. J., D. DeMartini, P. Dinnel and J. Siecke. 1976. The ecology of the gaper or horse clam, Tresus capax (Gould 1850) (Bivalvia: Mactridae), in Humboldt Bay, California. Calif. Fish and Game 62:41-64.
152. West, J. A. 1972. The life history of Petrocelis franciscana. Brit. Phycol. J. 7:299-308.
153. Whedon, W. F. 1936. Spawning habits of the mussel Mytilus californianus Conrad with notes on the possible relation to mussel poison I. Univ. Calif. Pub. Zool. 41:35-44.
154. Winzler and Kelly Consulting Engineers. 1977. A summary of knowledge of the central and northern California coastal zone and offshore areas. Vol. II. Biological conditions. Prepared for Bureau of Land Management, U.S. Dept. of the Interior, Contract AA550-CT6-52.
155. Young, R. T. 1942. Spawning season of the Californian mussel Mytilus californianus. Ecology 23:490-492.
156. Young, R. T. 1946. Spawning and setting season of the mussel, Mytilus californianus. Ecology 27:354-363.

BLM LIBRARY
RS 150A BLDG. 50
DENVER FEDERAL CENTER
P.O. BOX 25047
DENVER, CO 80225

R'S CARD

422 1980

tte L.
benthic marine
along the

OFFICE	DATE RETURNED			

(Continued on reverse)

TD 195 .P4 P422 1980
Vesco, Lynnette L.
Recovery of benthic marine
populations along the

BLM LIBRARY
RS 150A BLDG. 50
DENVER FEDERAL CENTER
P.O. BOX 25047
DENVER, CO 80225

